CORRECTION Open Access

Correction to: Laser-based molecular delivery and its applications in plant science



Dag Heinemann^{1,2,3*}, Miroslav Zabic^{1,2}, Mitsuhiro Terakawa⁴ and Jens Boch⁵

Correction to: Plant Methods (2022) 18:82

https://doi.org/10.1186/s13007-022-00908-9

In the original version of the article the wrong figure appeared as Fig. 1; Fig. 1 should have appeared as shown in this correction.

The original article [1] has been corrected.

The original article can be found online at https://doi.org/10.1186/s13007-022-00908-9.

¹ Hannover Centre for Optical Technologies, Leibniz University Hannover, Nienburger Str. 17, 30167 Hannover, Germany Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons locence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*}Correspondence: dag.heinemann@hot.uni-hannover.de

Heinemann et al. Plant Methods (2022) 18:104 Page 2 of 2




Fig. 1 Sketch of a typical photoinjection experiment using an inverted microscopic setup and a pulsed laser source. The laser beam is focused onto the sample using an inverted microscope setup. A single laser pulse or a train of pulses facilitates of the cellular membrane and possibly the cell wall. The exact physical process of photoporation depends on the applied laser parameters and will be discussed in the following section. Plasmolyzing the plant cell prior to photoinjection supports the molecular uptake

Author details

¹Hannover Centre for Optical Technologies, Leibniz University Hannover, Nienburger Str. 17, 30167 Hannover, Germany. ²Institute of Horticultural Production Systems, Leibniz University Hannover, Herrenhäuser Str. 2, 30419 Hannover, Germany. ³Cluster of Excellence PhoenixD, Leibniz University Hannover, Welfengarten 1, 30167 Hannover, Germany. ⁴Department of Electronics and Electrical Engineering, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan. ⁵Institute of Plant Genetics, Leibniz University Hannover, Herrenhäuser Str. 2, 30419 Hannover, Germany.

Published online: 25 August 2022

Reference

 Heinemann D, Zabic M, Terakawa M, Boch J. Laser-based molecular delivery and its applications in plant science. Plant Methods. 2022;18:82. https://doi.org/10.1186/s13007-022-00908-9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.