

Methodology

Open Access

A community resource for high-throughput quantitative RT-PCR analysis of transcription factor gene expression in *Medicago truncatula*

Klementina Kakar¹, Maren Wandrey¹, Tomasz Czechowski¹, Tanja Gaertner¹, Wolf-Rüdiger Scheible¹, Mark Stitt¹, Ivone Torres-Jerez³, Yongli Xiao², Julia C Redman², Hank C Wu², Foo Cheung², Christopher D Town² and Michael K Udvardi*^{1,3}

Address: ¹Max-Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam-Golm, Germany, ²The J. Craig Venter Institute, 9704 Medical Center Drive, Rockville, MD, 20850, USA and ³The Samuel Roberts Noble Foundation, 2510 Sam Noble Parkway, Ardmore, OK, 73401, USA

Email: Klementina Kakar - kakar@mpimp-golm.mpg.de; Maren Wandrey - wandrey@gfz-potsdam.de; Tomasz Czechowski - tc519@york.ac.uk; Tanja Gaertner - gaertner@mpimp-golm.mpg.de; Wolf-Rüdiger Scheible - scheible@mpimp-golm.mpg.de; Mark Stitt - stitt@mpimp-golm.mpg.de; Ivone Torres-Jerez - itjerez@noble.org; Yongli Xiao - yxiao@jcv.org; Julia C Redman - jredman@jcv.org; Hank C Wu - hwu@jcv.org; Foo Cheung - FCheung@jcv.org; Christopher D Town - cdtown@jcv.org; Michael K Udvardi* - mudvardi@noble.org

* Corresponding author

Published: 8 July 2008

Received: 15 April 2008

Plant Methods 2008, **4**:18 doi:10.1186/1746-4811-4-18

Accepted: 8 July 2008

This article is available from: <http://www.plantmethods.com/content/4/1/18>

© 2008 Kakar et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Medicago truncatula* is a model legume species that is currently the focus of an international genome sequencing effort. Although several different oligonucleotide and cDNA arrays have been produced for genome-wide transcript analysis of this species, intrinsic limitations in the sensitivity of hybridization-based technologies mean that transcripts of genes expressed at low-levels cannot be measured accurately with these tools. Amongst such genes are many encoding transcription factors (TFs), which are arguably the most important class of regulatory proteins. Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) is the most sensitive method currently available for transcript quantification, and one that can be scaled up to analyze transcripts of thousands of genes in parallel. Thus, qRT-PCR is an ideal method to tackle the problem of TF transcript quantification in *Medicago* and other plants.

Results: We established a bioinformatics pipeline to identify putative TF genes in *Medicago truncatula* and to design gene-specific oligonucleotide primers for qRT-PCR analysis of TF transcripts. We validated the efficacy and gene-specificity of over 1000 TF primer pairs and utilized these to identify sets of organ-enhanced TF genes that may play important roles in organ development or differentiation in this species. This community resource will be developed further as more genome sequence becomes available, with the ultimate goal of producing validated, gene-specific primers for all *Medicago* TF genes.

Conclusion: High-throughput qRT-PCR using a 384-well plate format enables rapid, flexible, and sensitive quantification of all predicted *Medicago* transcription factor mRNAs. This resource has been utilized recently by several groups in Europe, Australia, and the USA, and we expect that it will become the 'gold-standard' for TF transcript profiling in *Medicago truncatula*.

Background

Legumes are second only to grasses in agricultural importance [1]. They are a mainstay of sustainable agricultural systems because of their ability to reduce atmospheric nitrogen (N₂) to ammonia via a symbiosis with bacteria called rhizobia. This provides legumes and subsequent crops with a free and renewable source of nitrogen in lieu of expensive, environmentally-unfriendly fertilizers. Development and differentiation of root nodules, the organ that accommodates nitrogen-fixing rhizobia in legumes, is orchestrated by transcription factors [2-9]. Transcription factors are DNA-binding proteins that regulate the transcription of most, if not all genes [10]. As a result, TFs play central roles in all aspects of plant biology, including development and differentiation of organs and adaptive responses to changes in the environment [11]. Transcription factors as a whole are an important target of plant research because they are a key to understanding the regulation of important plant processes as well as potential tools to optimize these processes for agriculture.

The importance of TFs in plant biology is reflected by the fact that approximately 5% of all plant genes encode such proteins [10]. Thus, even species with relatively small genomes, such as *Arabidopsis thaliana* contain thousands of TF genes [10]. This presents a real challenge for systematic approaches to decipher the function of TF genes in plants. Classical, 'forward' genetics has uncovered the roles of perhaps a hundred TF genes in *Arabidopsis* [12] and far fewer in other species [11]. Reverse-genetic approaches, using T-DNA insertion mutants for instance [13], provide a means to decipher in a systematic and relatively rapid manner the function of TF genes/proteins, although gene-redundancy often stymies this enterprise [12]. Another stumbling-block is that phenotypes associated with non-redundant TFs may be subtle in nature.

Transcript profiling can help to uncover the functions of TF genes/proteins by revealing where and when in a plant TF genes are expressed. This information can help direct our attention to particular organs, developmental stages, or conditions under which aberrant phenotypes might become apparent in a TF mutant of interest.

Medicago truncatula is a model legume species that is currently the focus of an international genome sequencing effort [14]. Several generations of cDNA [15] and oligonucleotide arrays [16] have been developed for transcriptome analysis of *Medicago truncatula*, including most-recently an Affymetrix GeneChip that contains 51,000 probe-sets representing a large proportion of all the genes in this species [17]. While these tools now provide a means to measure the transcriptional output of a large proportion of genes in *Medicago*, inherent limitations in the sensitivity of hybridization-based technologies [18]

mean that transcripts of a substantial number of genes cannot be detected even when probes for these transcripts are present on the array/chip. Furthermore, expansion of arrays to encompass novel genes uncovered by genome sequencing is not a trivial task. An alternative to arrays that is 2–3 orders of magnitude more sensitive and more flexible in terms of expansion to encompass novel genes is quantitative reverse transcription-polymerase chain reaction (qRT-PCR). Platforms for qRT-PCR analysis of thousands of *Arabidopsis* and rice TF genes have been developed by us and others [19,20], and utilized to identify TF genes involved in *Arabidopsis* responses to nutrient stress and pathogen attack [21-24]. Here we describe a bioinformatics pipeline to identify putative TF genes in *Medicago truncatula* and to design gene-specific oligonucleotide primers for qRT-PCR analysis of all predicted TF transcripts. Over 1000 TF primer pairs were tested and used to identify sets of organ-enhanced TF genes that may play important roles in organ development or differentiation in this species.

Results and Discussion

Identification of putative transcription factors

TF protein families are generally defined by the type(s) of DNA-binding domain they contain and putative TF genes are often identified on the basis of DNA sequences that encode known DNA-binding domains [10,11,25,26]. We utilized this approach to identify putative TFs of *Medicago* amongst the set of proteins predicted from genomic sequence by the International *Medicago* Genome Annotation Group (IMGAG). Proteins of IMGAG release 1, which contained over 40,000 predicted proteins, were screened for the presence of known or presumed DNA-binding domains (Table 1), using INTERPRO [27]. *Medicago* proteins containing putative DNA binding domains and other domains associated with TFs were then used as query sequences in WU-BLASTX [28] which included searches of both the non-redundant DNA database of NCBI [29] and the well-curated protein database, UniProt [30] to check annotations of related proteins in support of tentative *Medicago* TF assignments. This process resulted in a list of 1045 putative TF genes (see Additional file 1). We utilized genomic sequences rather than the large collection of partial cDNA sequences present in Expressed Sequence Tag (EST) databases for *Medicago* as the starting point for TF gene discovery because protein sequences derived from genomic sequence are more complete and the set of IMGAG proteins essentially contains no redundancy. Although identification of the 'complete' set of *Medicago* TFs from IMGAG-annotated proteins will only be possible upon completion of genome sequencing, we expect little or no redundancy in the protein set targeted by our primer set. This approach avoids wasting money on redundant primer sets and re-organization of primers when redundancy is detected, both of which would have

Table 1: Classification of putative transcription factors of Medicago into families and sub-families

TF Family	No. of Genes	Characteristic Domain (InterPro No.)	Domain Function	Domain Description
MYB/HD-like	76	IPR001005, IPR009057	D	Myb, DNA-binding; homeodomain like
MYB	58	IPR001005	D	Myb, DNA-binding
C ₂ H ₂ (Zn)	64	IPR007087	NA	Zn-finger, C ₂ H ₂ type
AP2/EREBP	55	IPR001471	D	Pathogenesis-related transcriptional factor and ethylene response factor
BHLH	49	IPR001092	D	Basic helix-loop-helix dimerisation region bHLH
HD-like	50	IPR009057	D	Homeodomain like
HD family		IPR001356	D	Homeobox
HD	25			
HD-ZIP	5	IPR006712	P	HD-ZIP protein, N terminus
HD-PHD finger	2	IPR001965	P	Zn-finger like, PHD-finger
MADS	48	IPR002100	D	TF, MADS-box
BZIP	41	IPR004827	D	Basic Leu zipper (bZIP) TF
PHD	34	IPR001965	P	Zn-finger like, PHD-finger
WRKY family		IPR003657	D	DNA-binding WRKY
WRKY	29			
LLR WRKY	1	IPR001611		Leu-rich repeat
ABI3/VP1	29	IPR003340	D	TF B3
NAC	29	IPR003441	D	No apical meristem (NAM) protein
C ₃ H-type I (Zn)	27	IPR000571	D	Zn-finger, C-x8-C-x5-C-x3-H type
ARF	23	IPR003340, IPR010525, IPR011525	D	
JUMONJI	20	IPR003347	D	TF jumonji, jmjC
GRAS	19	IPR005202	P	GRAS TF
HMG	15	IPR000637	D	HMG-I and HMG-Y, DNA binding
AS2	14	IPR004883	P	Lateral organ boundaries
C ₂ C ₂ (Zn)				
Dof	13	IPR003851	D	Zn-finger, Dof type
GATA	7	IPR000679	D	Zn-finger, GATA type
CO-like	6	IPR000315	D	Zn-finger, B-box
YABBY	5	IPR006780	D	YABBY protein
CCAAT-HAP3 type	12	IPR003958	D	TF CBF/NF-Y/archaeal histone
GRF	8	IPR010666	D	Zn-finger, GRF type
SBP	8	IPR004333	D	SBP
EIL	7	IPR006957	D	Ethylene insensitive 3
LIM	7	IPR001781	P	Zn-binding protein, LIM
SNF2	6	IPR000330	D	SNF2 family N-terminal domain
E2F/DP	5	IPR003316	D	TF E2F/dimerisation partner (TDP)
TCP	5	IPR005333	D	TCP TF
FHA	5	IPR000253	D	Forkhead-associated
ARID	4	IPR001606	D	AT-rich interaction region
HSF	4	IPR000232	D	Heat shock factor (HSF)-type, DNA binding
AUX/IAA	3	IPR003311	D	AUX/IAA protein
SRS	3	IPR006510	D	Zn-finger, LRP1 type
TUB	3	IPR000007	D	Tubby
ZIM	3	IPR010399	D	ZIM
DDT	3	IPR004022	D	DDT
ZF-HD	2	IPR006455	D	Homeobox domain, ZF-HD class
MBF1	2	IPR001387	D	Helix-turn-helix type 3
SIFa-like	2	IPR006779	D	DNA binding protein SIFA
CAMTA	2	IPR005559	D	CG-I
LFY	1	IPR002910	D	Floricaula/leafy protein
Nin-like	1	IPR003035	D	Plant regulator RWP-RK
TAZ	1	IPR000197	P	Zn-finger, TAZ-type

Potentially novel plant TFs and transcriptional regulators

CCHC (Zn)	112	IPR001878	NA	Zn-finger, CCHC-type
RR	16	IPR001789, IPR011006	RD	Response regulator receiver

Table 1: Classification of putative transcription factors of Medicago into families and sub-families (Continued)

DHHC (Zn)	14	IPR001594	D or P	Zn-finger, DHHC-type
HTH				
	FIS	IPR002197	D	Helix-turn-helix, Fis-type
	AraC	IPR000005	D	Helix-turn-helix, AraC type
BTB/POZ	7	IPR000210	P	BTB
TTF-type (Zn)	6	IPR006580	D	Zn-finger, TTF-type
BD	6	IPR001487	P	Bromodomain
Lambda-DB	3	IPR010982	D	Lambda_DNA_bd
TrpR	3	IPR010921	D	Trp repressor/replication initiator
TPR	3	IPR001440	P	Tetratricopeptide TPR_I
KRAB-box	2	IPR001909	P	KRAB box
NRs	2	IPR008946	LBD	Steroid nuclear receptor, ligand binding
R3H	2	IPR001374	NA	Single-stranded nucleic acid binding R3H
YEATS	2	IPR005033	TA	YEATS
U1-type (Zn)	2	IPR003604	NA	Zn-finger, U1-type
A20-like	2	IPR002653	P	Zn-finger, A20-type
Euk_TF	1	IPR008917	D	Euk_TF, DNA binding
NGN	1	IPR006645	D	NGN
p53-like	1	IPR008967	D	p53-like TF, DNA binding
SSB protein	1	IPR011344	D	Single-strand binding protein
ssDB TR	1	IPR009044	D	Single-strand DNA binding transcriptional regulator
TCoAp15	1	IPR003173	D	Transcriptional coactivator p15
BED-type (Zn)	1	IPR003656	D	Zn-finger, BED-type predicted
TCoA	1	IPR009255	TA	Transcriptional coactivation
Tc/PD	1	IPR001533	TA	Transcriptional coactivator

IMGAG (International Medicago Gene Annotation Group)-proteins were classified as putative TFs if they contained characteristic DNA-binding or other characteristic TF domains and if annotations of matching proteins obtained by BLAST searches were consistent with such a classification. TF families previously identified in plants are presented in the first part of the table while potentially novel plant TF families, which were identified by the presence of domains associated with TFs and other transcriptional regulators outside the plant kingdom, are presented in the latter part of the table. D = DNA binding domain; P = protein-protein interaction domain; NA = nucleic acid (DNA and RNA) binding domain; RD = receiver domain; LBD = ligand binding; transcriptional co-activator. Plant-specific TF families and sub-families are indicated in bold (according to [12])

been inevitable if we chose to use ESTs in addition to genomic sequences to identify Medicago TFs.

PCR primer design

To ensure maximum specificity and efficiency during PCR amplification of TF cDNA under a standard set of reaction conditions, a stringent set of criteria was used for primer design. This included predicted melting temperatures (T_m) of 58°C to 61°C, limited self-complementarity and poly-X, and PCR amplicon lengths of 100–150 base pairs (bp). Secondary hits were minimized by aligning primer candidates to all known Medicago sequences via WU-BLAST [28] and eliminating primer pairs with multiple potential hits.

PCR primer testing: gene-specificity and amplification efficiency

PCR primers were tested on Medicago cDNA free of genomic DNA contamination as follows. First, total RNA was extracted from various organs using Trizol reagent (Invitrogen GmbH, Karlsruhe, Germany), which yielded high quality RNA as judged by gel electrophoresis and by Agilent 2100 BioAnalyser using RNA 6000 Nano Chips (Agilent Technologies, Waldbronn, Germany). Typical RNA yields ranged from 0.5–1.0 µg RNA/mg fresh mass for nodules and leaves, respectively. Isolated RNA was

treated with DNase I (Ambion, product number 1907) to remove all contaminating genomic DNA, and this was always confirmed by PCR using primers to non-coding regions of the *Ubiquitin* gene (TC102473; AC137828-19.4). After inactivation of DNase I, RNA was reverse transcribed using SuperScript III reverse transcriptase (Invitrogen GmbH, Karlsruhe, Germany) and oligo-dT₁₂₋₁₈ to prime the reaction.

Specificity of PCR primers was assessed in three ways: by melting curve analysis of PCR reaction products; by separating the products of all reactions via electrophoresis in 3% agarose gels; and by sequencing a sub-set of PCR reaction products (Figure 1). 94.5% (998/1045) of primer pairs gave unique PCR products of the expected size. Only 3.3% (34/1045) of primer pairs yielded no product and 2.2% (23/1045) gave non-specific products. (see Additional file 1). Sequencing was performed on 178 randomly-chosen PCR products amplified from a 1:1 mixture of leaf and root cDNA. In the vast majority of cases (92.7% or 165/178), the sequence of the PCR product was identical to that of the intended target gene. In 5.1% of cases, the amplicon sequence matched multiple related genes, including the target gene, while in only 2.2% of cases the amplicon sequence did not match the target gene sequence.




Figure 1
Specificity of transcription factor PCR primers. Specificity was confirmed by dissociation curves with a single peak (A) while double peaks (B) indicated off-target amplification. The derivative of fluorescence intensity is shown on the y-axis. Separation of PCR products on 3% (v/w) agarose gels following electrophoresis (C) confirmed the presence of unique amplicons of the expected size for most reactions. Few reactions yielded no products (indicated by arrow).

Ideally, PCR results in an exact doubling of the amount of dsDNA after each temperature cycle. In practice, however, this is generally not the case because the reactions are less than 100% efficient. Primer sequences can affect PCR efficiency, so we determined the efficiency of each TF primer pair from amplification plots, using LinRegPCR software [31]. First, the correlation coefficient derived from linear regression analysis of each amplification plot (e.g. see Figure 2) was used to assess the 'quality' of each reaction, and all reactions with an $R^2 < 0.990$ were excluded from further analysis (10.6% of reactions). Next, average PCR efficiencies (E) were computed for each individual primer pair across all analyzed samples. 53.4% (558 TF genes) displayed PCR efficiencies greater than 0.80, while 39.7%

(415 TF genes) had efficiencies between 0.51–0.80. Only 2.6% (27 TF genes) had mean E values below 0.4; these all yielded $R^2 < 0.99$ in LinRegPCR analysis and mostly represented reactions that lacked detectable fragment amplification ($C_T > 40$) or that generated unspecific PCR products (Figure 2; see Additional file 1). A similar range of PCR efficiencies were determined for Arabidopsis and rice TF primers previously [19,20].

Selection of reference genes

Reference genes with stable expression/transcript levels throughout development and in the face of environmental challenge are crucial for the normalization of expression data of other genes. Potentially useful reference genes




Figure 2
Amplification efficiency of transcription factor-specific primer pairs. Typical real-time RT-PCR amplification plots of 384 TF genes (left) and distribution of PCR efficiencies for all 1045 TF primer pairs (right).

were chosen based on published data for *Medicago* (e.g. *Msc27* [32]) and *Arabidopsis thaliana* [33]. The closest *Medicago* homologues of *Arabidopsis* genes were identified by BlastN [29]. Gene-specific primer pairs for the genes encoding elongation factor 1 α (EST317575), glyceraldehyde-3-phosphate dehydrogenase (MtC00030_GC; CT573421_3.4), β -tubulin (TC106341), Pentatricopeptide repeat protein (TC96273), actin2 (TC107326; AC137836_27.5), Ubiquitin (TC102473; AC137828-19.4), Helicase (CB892427), and the genes *PDF2* (TC107161), *UPL7* (TC111218), *PTB* (TC111751), *UBC* (AW686873), *bHLH* (CX538576), and *UBC9* (TC106312) were designed using the criteria described above (Table 2). The specificity of PCR primers was tested using 18 first-strand cDNAs from six different organs of *Medicago* (three biological replicates each). All primer pairs produced a single PCR product of the expected size, as shown by gel electrophoresis and unique dissociation curves generated by the PCR machine after 40 cycles (Figure 3). To determine which reference genes were best suited for transcript normalisation, we used the software geNORM [34], which uses pair-wise comparison and geometric averaging across a matrix of biological samples to

determine gene expression stability (M; [35]). The genes *PDF2*, *PPRep*, *Ubiquitin*, and *PTB* had the lowest M (greatest transcript stability) and, therefore, were judged to be the best reference genes for this diverse set of developmental samples (Figure 4; Table 2).

Identification of organ-enhanced TFs of *Medicago*

To get an overview of TF gene expression in *Medicago truncatula* and to identify TFs induced in specific organs, we used the real-time RT-PCR platform described above. Transcript profiling was performed on six different organs of *Medicago* (leaves, stems, flowers, pods, roots, and nodules) with three independent biological replicates for each (see Additional file 2). The fraction of genes for which transcripts were detected within 40 cycles ranged from 77.2% in leaves to 90.8% in pods. Transcripts from nearly all putative TF genes (96.8% or 1011/1045) were detected in at least one organ. Genes were called detected if they were expressed in at least two biological replicates with a $C_T < 40$. Approximately half of all TF genes exhibited differential expression during plant development, based on significant differences ($p \leq 0.05$) in transcript levels between organs. Few TF genes (1.19% or 12/1011) were

Table 2: Medicago reference genes and primers for qRT-PCR

Gene Name	TC	Accession Number	Forward/Reverse Primer (5'-3')	PCR Product Size (bp)	PCR efficiency (E)	R ²
β Tubulin	TC106341	N	TTTGCTCCTTACATCCCGTG / GCAGCACACATCATGTTTTTGG	100	1.08	1.0 0
PPRrep	TC96273	N	GGAAACTGGAGGATGCACGTA / ACAAGCCCTCGACACAAAACC	100	0.93	1.0 0
PDF 2	TC107161	N	GTGTTTTGCTTCCGCCGTT / CCAAATCTTGCTCCCTCATCTG	100	0.99	1.0 0
bHLH	CX538576	N	TAGCGAGTACCATGATGCCAGA / GCGCCTCTTTTGTTCAGC	100	0.89	1.0 0
UBC	AW686873	N	CTGACAGCCCACTGAATTGTGA / TTTTGGCATTGCTGCAAGC	100	0.96	1.0 0
PTB	TC111751	N	CGCCTTGTCAGCATTGATGTC / TGAACCAAGTGCCTGGAATCCT	100	0.85	1.0 0
Ubiquitin	TC102473	AC137828_19.4	GCAGATAGACACGCTGGGA / AACTCTTGGGCAGGCAATAA	100	0.95	1.0 0
UBC9	TC106312	AC137602_2.4	GGTTGATTGCTCTTCTCCTCCC / AAGTGATTGCTCGTCCAACCC	100	1.13	0.9 9
Helicase	CB892427	N	GTACGAGGTGGTGCTCTTGAA / GCAACCGAAAATTGCACCATAC	100	0.91	1.0 0
ELF1α	EST317575	N	GACAAGCGTGTGATCGAGAGATT / TTTCACGCTCAGCCTTAAGCT	100	0.68	0.9 8
UPL7	TC111218	N	CCAGTTGTTCTCGTGGTCCATT / CCTCCAATTGTCGCCAAA	100	0.93	1.0 0
GAPDH	MtC00030_GC	CT573421_3.4	TGCCCTCTGATTCTCCTTG / TGCCCTCTGATTCTCCTTG	100	1.04	0.9 9
Actin2	TC107326	AC137836_27.5	TCAATGTGCCTGCCATGTATGT / ACTCACACCGTCACCAGAATCC	100	1.12	0.9 9
MSC27	X63872 (<i>M. sativa</i>)	N	GTTGAAGTAGACATTGGTGCTAACG / AGCTGAGTCATCAACCCCTCAT	100	0.76	0.9 9

Mean PCR efficiency (E) was determined from three biological replicates of each of six organs, using LinRegPCR [31], which also yielded mean R². N, no corresponding GenBank accession number.

expressed exclusively in vegetative organs (leaves, stems, roots or nodules), and even fewer (0.5% or 5/1011) were expressed only in reproductive organs (flowers or pods). A relatively small number of TF genes exhibited greater than ten-fold ratios in expression level in one organ compared to any other organ (Table 3). For comparison, we have included gene expression ratios derived from Affymetrix array data from the same RNA samples. While there is reasonable qualitative agreement between gene expression ratios obtained using the two methods, the lack of quantitative agreement is likely due to the limited sensitivity and low signal to noise ratio near the detection limit of Affymetrix arrays [19]. The genes listed in Table 3 may control development and/or differentiation in *Medicago* and are interesting targets for future research.

Conclusion

We have established a flexible platform for high-throughput qRT-PCR analysis of *Medicago* TF gene expression that is based on gene-specific primers arrayed in 384-well plates and SYBR® Green detection of gene-specific PCR amplicons. Currently, the platform has primer pairs for 1045 TF genes and we have plans to extend this to all pre-

dicted *Medicago* TF genes as genome sequencing progresses. At this stage, the resource has been utilized by several groups in Europe, Australia, and the USA, and we expect it will become the 'gold-standard' for TF transcript profiling in *Medicago truncatula*.

Methods

Plant material and growth conditions

Medicago truncatula cv. Jemalong, line A17 wild type plants were vernalized for 3 days in the dark at 4°C on sterile, wet filter paper. Germinated seedlings were transferred to pots containing Turface (BWI Texarcana, Texarcana, TX). Plants were grown in growth chambers under a 16 h day and 8 h night regime, at 200 μE light intensity, 24°C and 40% relative humidity.

Vegetative organs (leaves, stems, roots, and nodules) were harvested 28 days after planting. Leaf material did not include petioles and stems did not include buds. Roots consisted of the entire root system with laterals. Several plants grown at the same time were pooled for each of the three biological replicates. Biological replicates were planted on separate days. Nodules were harvested from




Figure 3
Specificity, efficiency, and reproducibility of PCR primers designed to amplify reference gene transcripts. Specificity of primers was confirmed by the presence of unique amplicons of the expected size following electrophoresis on 3% (v/w) agarose gels (A) and by dissociation curves with a single peak (B to D). Typical real-time RT-PCR amplification plots of three reference gene transcripts (E to G).




Figure 4
Ranking of 8 reference genes in *M. truncatula*. Transcript levels of all 8 genes were measured by qRT-PCR, using 18 independent cDNA preparations from six different organs with three replicate measurements of each cDNA preparation. A low value for the average expression stability M, as calculated by geNORM software, indicates more stable expression throughout the various organs.

plants inoculated with *Sinorhizobium meliloti* strain 1021 one and seven days after sowing. Reproductive organs were harvested from plants that were vernalized for two weeks to decrease the time to flowering. Flowers were harvested on the day of opening. Pods were harvested from 1 to 21 days after the appearance of the floral bud to cover a wide range of developmental stages. Harvested plant material was frozen in liquid nitrogen before storage at -80°C.

RNA isolation and cDNA synthesis

Total RNA was extracted using Trizol reagent [36], following the manufacturer's instructions (Invitrogen GmbH, Karlsruhe, Germany). RNA was quantified using a NanoDrop Spectrophotometer ND-100 (NanoDrop Technologies, Wilmington, DE). Sixty µg of total RNA were digested with RNase free DNase1 (Ambion Inc., Houston, TX), according to manufacturer's protocol. RNA integrity was checked using an Agilent 2100 BioAnalyser and RNA 6000 Nano Chips (Agilent Technologies, Waldbronn, Germany), and by electrophoresis on a 3% (v/w) agarose gel before and after DNase I treatment. The absence of contaminating genomic DNA after DNase I treatment was verified by PCR analysis, using primer pairs designed to amplify a 107 bp genomic fragment of the control gene, *Ubiquitin* (TC102473intronF, 5'-GTCCTCTAAGGTTAATGAACCGG-3'; TC102473intronR, 5'-GAAAGACACAGCCAAGTTGCAC-3').

First-strand complementary DNA was synthesized by priming with oligo-dT₁₂₋₁₈ (Qiagen, Hilden, Germany), using SuperScript III reverse transcriptase (Invitrogen GmbH, Karlsruhe, Germany) following the instructions of the provider. To assess cDNA synthesis efficiency, qPCR was used to amplify segments in the 5' and 3' regions of *Ubiquitin* cDNA approx. 1600 and 400 bp from the 3'-end, respectively (primers: TC102473_5'F, 5'-TTGGAGACG-GATTCCATTGCT-3'; TC102473_5'R, 5'-GCCAATTCCT-TCCCTTCGAA-3; TC102473_3'F, 5'-GGCCCTAGAACATTTCTGTGG-3'; and TC102473_3'R, 5'-TTGGCAACCAAATGTCCCC-3'). If ΔCt (Ct₃-Ct₅) < 2, then cDNA synthesis efficiency was judged to be satisfactory, and the cDNA was considered suitable for qRT-PCR analysis.

PCR primer design

The primer design pipeline was implemented in object-oriented PERL modules supported by a MySQL relational database. Primers iterated through three phases before approval: design, specificity, and selection.

The design phase interrogated TF genes with a sliding window 250 bp across that stepped 50 bp along the entire target sequence, generating primer candidates at each window. Experimental conditions, as outlined in the Results section above, were enforced by the following MIT Primer3 parameters: PRIMER_MIN_TM 58, PRIMER_OPT_TM 60, PRIMER_MAX_TM 61, PRIMER_SELF_ANY 6, PRIMER_SELF_END 2, PRIMER_MAX_POLY_X 3, and PRIMER_PRODUCT_SIZE_RANGE '100-150' [37]. The specificity phase aligned primer candidates via WU-Blast to a database of all known Medicago sequences. The selection phase sorted primer candidates by the number of possible secondary hits, self-complementarity, and poly-X characteristics. Secondary hits were defined as specificity alignments that contained at least one of the terminal ends of the primer and achieved 80% or greater identity over the length of the primer. The sequences of each primer pair are given in Supplementary Material (see Additional file 1).

Real-time PCR conditions and analysis

PCR reactions were carried out in an ABI PRISM® 7900 HT Sequence Detection System (Applied Biosystems, Foster City, CA, USA). SYBR® Green was used to quantify dsDNA synthesis. Reactions (5 µl total volume) were performed in an optical 384-well plate containing 2.5 µl 2 × SYBR® Green Power Master Mix reagent (Applied Biosystems, Warrington, UK), 5 ng cDNA and 200 nM of each gene-specific primer. Primer pairs were aliquoted using a pipetting robot (Evolution P3 liquid handling system, Perkin Elmer, MA, USA) to minimize pipetting errors. cDNA was aliquoted as a master mix of cDNA and 2 × SYBR® Green

Table 3: Organ-enhanced TF genes

A Root-enhanced TF genes identified by real-time RT-PCR

Accession Number	Real-time RT-PCR Expression Ratio					Affymetrix Expression Ratio					Affymetrix Chip ID
	R/L	R/S	R/F	R/P	R/N	R/L	R/S	R/F	R/P	R/N	
ACI40721_12.1	121.5	320.1	1623.4	2676.5	64.4	161.2 ^b	144.5 ^b	219.2 ^b	158.1 ^b	99.9	Mtr.50075.1.SI_s_at
ACI35101_25.1	312.2	438.0	291.6	787.0	61.3	n	n	n	n	n	n
ACI40721_14.1	1204.9	751.2	956.6	6296.6	32.6	n	n	n	n	n	n
ACI40031_3.1	275.4	1157.3	923.4	258916.9 ^a	31.1	0.4	0.7	0.8	0.9	2.2	Mtr.47227.1.SI_s_at
ACI40721_13.1	39.9 ^a	179.8	614.8	935.1 ^a	26.5	35.9	48.1 ^b	41.2	44.2 ^b	39.5	Mtr.50074.1.SI_at
ACI40031_7.1	360.2 ^a	830.6	1542 ^a	6868.8	19.7	0.7 ^b	1 ^b	1.1 ^b	1 ^b	0.9 ^b	Mtr.47229.1.SI_at
ACI46574_6.1	1095.7	11.9^a	48.8	51.8	88.8	1.2 ^b	1.1 ^b	1.3 ^b	1.1 ^b	1.2 ^b	Mtr.40781.1.SI_s_at
ACI25478_13.7	1117.4	7008.0	3264.9 ^a	7256.6	11.6	211.5 ^b	248.8 ^b	296.4 ^b	228.7 ^b	45.3	Mtr.15416.1.SI_at
ACI25478_7.2	1052.3	503.3	938.4 ^a	2754.5	11.2	n	n	n	n	n	n
ACI22726_21.111	50.0	48.0	20943.9	2658.6	10.4	74.3 ^b	76.4 ^b	72.8 ^b	82.8 ^b	26.6	Mtr.15568.1.SI_s_at

B Nodule-enhanced TF genes identified by real-time RT-PCR

Accession Number	Real-time RT-PCR Expression Ratio					Affymetrix Expression Ratio					Affymetrix Chip ID
	N/L	N/S	N/F	N/P	N/R	N/L	N/S	N/F	N/P	N/R	
ACI48816_3.2	9966.7	1826.2	6262.9	21709.1	662.7	434.6 ^b	516.8 ^b	455 ^b	521.7 ^b	551.4 ^b	Mtr.14503.1.SI_at
ACI47774_3.2	55.2^a	466.7	726.3	881.2	498.5	17.7 ^b	16 ^b	17.8 ^b	19.1 ^b	16.1 ^b	Mtr.19554.1.SI_at
ACI38056_33.241	43.9	93.2	18.3	17.3	13.1	0.9 ^b	0.8 ^b	0.8 ^b	1.1 ^b	1.1 ^b	Mtr.17993.1.SI_at
ACI24214_39.2	121.8	92.1	44.8	64.9	10.5	n	n	n	n	n	n

C Pod-enhanced TF genes identified by real-time RT-PCR

Accession Number	Real-time RT-PCR Expression Ratio					Affymetrix Expression Ratio					Affymetrix Chip ID
	P/L	P/S	P/F	P/N	P/R	P/L	P/S	P/F	P/N	P/R	
ACI43340_4.7	154.1	44.8	173.3	180.3 ^a	407.3	1.6 ^b	1.8 ^b	1.7 ^b	1.7 ^b	1.7 ^b	Mtr.17931.1.SI_at

D Flower-enhanced TF genes identified by real-time RT-PCR

Accession Number	Real-time RT-PCR Expression Ratio					Affymetrix Expression Ratio					Affymetrix Chip ID
	F/L	F/S	F/P	F/N	F/R	F/L	F/S	F/P	F/N	F/R	
ACI29092_13.1	40.3	383.3	65.1	1683.9	318.1	24.2	47.1 ^b	38.2	45.4 ^b	55.1 ^b	Mtr.16432.1.SI_at
ACI48485_10.1	476.2	203.7	27.2	174.6	44.5	36.5 ^b	32.9 ^b	22.0	34.7 ^b	29.7 ^b	Mtr.20392.1.SI_at
ACI40915_20.1	169.5	101.1	18.6	481.0	178.7	1.4 ^b	1.4 ^b	1.3 ^b	1.3 ^b	1.6 ^b	Mtr.51688.1.SI_at
ACI141107_50.2	38.2	33.0	17.9	114.0	97.4	n	n	n	n	n	n
ACI44731_15.21	64.4	44.3	33.6	61.5	16.3	0.6 ^b	0.6 ^b	0.8 ^b	0.8 ^b	0.8 ^b	Mtr.19093.1.SI_at
ACI50978_12.1	235.3	198.5	48.4	14.0	16.9	n	n	n	n	n	n
ACI41107.5.61	261.1	127.8	13.3	96.4	71.5	0.9 ^b	0.9 ^b	0.8 ^b	0.1 ^b	1.1 ^b	Mtr.51651.1.SI_at
ACI57472_19.1	60.2 ^a	92.1	13.2	61.5 ^a	113.6	n	n	n	n	n	n
ACI48527_19.141	188.4	42.1	17.3	26.5	13.0	n	n	n	n	n	n
ACI44726_6.1	254.7	55.2	10.9	112.7 ^a	754.7	59.2	73.7 ^b	7.6	89.2	97.9 ^b	Mtr.19024.1.SI_at
ACI57488_16.1	7053.2 ^a	141.1	18.7	81.8	10.7	n	n	n	n	n	n
ACI23899_15.181	10.2	92.2	19.1	62.8	24.9	0.9 ^b	0.5 ^b	0.8 ^b	0.1 ^b	0.8 ^b	Mtr.52015.1.SI_at

A TF gene was considered organ-enhanced if transcript levels for that gene were more than 10-fold higher in one organ than in any other organ. Transcript ratios were calculated using the mean of three biological replicates for each organ. Data from qRT-PCR are compared with data for the same RNA samples obtained from Affymetrix Gene chips [38]. Affymetrix data were normalized using the Robust Multiarray Average (RMA) method, as described by [38], prior to calculation of ratios. Data in bold represent the lowest transcript ratio of the corresponding gene across all organs. n = not present on Affymetrix chip; a = Ct > 40 in two or three biological replicates of denominator organ; b = transcript called 'absent' by Affymetrix software in two or three biological replicates of denominator organ. L = leaf; S = stem; P = pod; F = flower; R = root; N = nodule.

reagent, using an electronic Eppendorf multipipette. Reaction plates were sealed with a transparent adhesive cover before proceeding (Applied Biosystems, Foster City, CA, USA). All templates were amplified using the following standard PCR protocol: 50 °C for 2 min; 95 °C for 10 min; 40 cycles of 95 °C for 15 sec and 60 °C for 1 min, and SYBR® Green fluorescence was measured continuously. Melting curves were generated after 40 cycles by heating the sample up to 95 °C for 15 sec followed by cooling down to 60 °C for 15 s and heating the samples to 95 °C for 15 sec.

Data analysis was performed with the SDS 2.2.1 software (Applied Biosystems). To determine the threshold cycle value (C_T) for each PCR reaction, the threshold (ΔR_n) was set within the logarithmic amplification phase. All amplification plots were analyzed with an ΔR_n of 0.2. PCR efficiency (E) was estimated using LinReg software with data obtained from the exponential phase of each individual amplification plot and the equation $(1+E) = 10^{\text{slope}}$ [31]. To compare data from different PCR runs and different cDNA samples, C_T values were normalized against the geometric mean of four reference genes (*Ubiquitin*, *PPRep*, *PDF2*, and *PTB*), whose transcript levels were most stable across the biological samples analyzed. The average of the geometric mean of these four genes for all 18 samples was $C_{T-21.23} \pm SD1.15$. For normalization, the mean reference gene C_T value was subtracted from the C_T value of the TF gene of interest, yielding a ΔC_T value. The expression ratios for the identification of organ-enhanced genes were obtained using the following formula on all 30 organ combinations: $(1 + E)^{\Delta\Delta C_T}$, where $\Delta\Delta C_T$ was calculated by ΔC_{TA} minus ΔC_{TB} , A and B are averages of three biological replicates of the two organs being compared, and E is the PCR efficiency. Dissociation curves were analysed using SDS 2.2.1 software (Applied Biosystems). RT-PCR products were resolved on 3% (w/v) agarose gels (LE Agarose, Biozym, Oldendorf, Germany) run at 4 V cm^{-1} in TAE Tris-Acetate-EDTA buffer, along with a 200-bp DNA-standard ladder (Promega GmbH). A subset of 178 RT-PCR products was sequenced at the JC Venter Institute (Rockville, MD, USA).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KK performed the experimental work and helped draft the manuscript. W-RS, TC, and MS helped to conceive the project and provided practical advice. MW, FC, and CDT

carried out bioinformatic analysis of Medicago TFs, and HW designed gene-specific primers. JCR and YX were responsible for the sequencing and analysis of the PCR products. IT carried out qRT-PCR and TG performed statistical analyses. MU designed and coordinated the project and wrote the manuscript.

Additional material

Additional file 1

Complete list of TF genes, primer sequences and corresponding PCR efficiencies. The TF primer platform was established based on the data for the gene models according to SA, specific amplification; NA, no amplification; NS, non-specific amplification. RT-PCR products of primer sequences indicated in bold were sequenced.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1746-4811-4-18-S1.xls>]

Additional file 2

Complete list of TF genes, gene families and experimental data. Shown are the Medicago gene Accession Numbers (TIGR) in ascending order, the TC number if available, as well as the transcription factor family and the subfamily names. The next columns show experimental results for the real-time RT-PCR reactions performed on six different organs of Medicago in three independent biological replicates, as explained in the manuscript. CT = not normalized CT value, ΔCT = CT value normalized against the geometric mean of 4 house keeping genes; $\Delta\Delta CT$ = $\text{power}(PCR_{\text{eff}}; \Delta CT)$; $\log_2 \Delta\Delta CT$ = logarithmus of $\Delta\Delta CT$.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1746-4811-4-18-S2.xls>]

Acknowledgements

This work was supported in part by the European Commission FP6 Framework Programme Grain Legume Integrated Project (FOOD-CT-2004-506223), the Max Planck Society, the National Research Initiative (NRI) Plant Genome Program of the USDA Cooperative State Research, Education and Extension Service (CSREES), and the Samuel Roberts Noble Foundation.

References

- Graham PH, Vance CP: **Legumes: importance and constraints to greater use.** *Plant Physiology* 2003, **131**:872-877.
- Schauser L, Roussis A, Stiller J, Stougaard J: **A plant regulator controlling development of symbiotic root nodules.** *Nature* 1999, **402(6758)**:191-195.
- Frugier F, Poirier S, Satiat-Jeunemaitre B, Kondorosi A, Crespi M: **A Kruppel-like zinc finger protein is involved in nitrogen-fixing root nodule organogenesis.** *Genes Dev* 2000, **14(4)**:475-482.
- Nishimura R, Ohmori M, Fujita H, Kawaguchi M: **A Lotus basic leucine zipper protein with a RING-finger motif negatively regulates the developmental program of nodulation.** *Proceedings of the National Academy of Sciences of the United States of America* 2002, **99(23)**:15206-15210.
- Gronlund M, Gustafsen C, Roussis A, Jensen D, Nielsen LP, Marcker KA, Jensen EO: **The Lotus japonicus ndx gene family is involved in nodule function and maintenance.** *Plant Molecular Biology* 2003, **52(2)**:303-316.
- Kaló P, Gleason C, Edwards A, Marsh J, Mitra RM, Hirsch S, Jakab J, Sims S, Long SR, Rogers J, Kiss GB, Downie JA, Oldroyd GED: **Nodulation signaling in legumes requires NSP2, a member of the**

- GRAS family of transcriptional regulators.** *Science* 2005, **308(5729)**:1786-1789.
7. Smit P, Raedts J, Portyanko V, Debelle F, Gough C, Bisseling T, Geurts R: **NSPI of the GRAS protein family is essential for rhizobial Nod factor-induced transcription.** *Science* 2005, **308(5729)**:1789-1791.
 8. Combier JP, Frugier F, de Billy F, Boualem A, El-Yahyaoui F, Moreau S, Vernie T, Ott T, Gamas P, Crespi M, Niebel A: **MtHAP2-I is a key transcriptional regulator of symbiotic nodule development regulated by microRNA169 in *Medicago truncatula*.** *Genes & Development* 2006, **20(22)**:3084-3088.
 9. Middleton PH, Jakab J, Penmetsa RV, Starker CG, Doll J, Kaló P, Prabhur R, Marsh JF, Mitra RM, Kereszt A, Dudas B, VandenBosch K, Long SR, Cook DR, Kiss GB, Oldroyd GE: **An ERF transcription factor in *Medicago truncatula* that is essential for Nod factor signal transduction.** *Plant Cell* 2007, **19(4)**:1221-1234.
 10. Riechmann JL, Heard J, Martin G, Reuber L, Jiang CZ, Keddie J, Adam L, Pineda O, Ratcliffe OJ, Samaha RR, Creelman R, Pilgrim M, Broun P, Zhang JZ, Ghandehari D, Sherman BK, Yu GL: **Arabidopsis transcription factors: Genome-wide comparative analysis among eukaryotes.** *Science* 2000, **290(5499)**:2105-2110.
 11. Udvardi MK, Kakar K, Wandrey M, Montanari O, Murray J, Andriankaja A, Zhang JY, Benedetto V, Hofer JMI, Chueng F, Town CD: **Legume Transcription Factors: Global Regulators of Plant Development and Response to the Environment.** *Plant Physiol* 2007, **144(2)**:538-549.
 12. Riechmann JL: **Transcriptional Regulation: a Genomic Overview.** In *The Arabidopsis Book* Edited by: Somerville CR, Meyerowitz EM. The American Society of Plant Biologists, Rockville; 2002.
 13. Alonso JM, Ecker JR: **Moving forward in reverse: genetic technologies to enable genome-wide phenomic screens in Arabidopsis.** *Nat Rev Genet* 2006, **7(7)**:524-536.
 14. Young ND, Cannon SB, Sato S, Kim D, Cook DR, Town CD, Roe BA, Tabata S: **Sequencing the genespaces of *Medicago truncatula* and *Lotus japonicus*.** *Plant Physiol* 2005, **137(4)**:1174-1181.
 15. El-Yahyaoui F, Küster H, Amor BB, Hohnjec N, Pühler A, Becker A, Gouzy J, Vernie T, Gough C, Niebel A, Godiard L, Gamas P: **Expression profiling in *Medicago truncatula* identifies more than 750 genes differentially expressed during nodulation, including many potential regulators of the symbiotic program.** *Plant Physiology* 2004, **136(2)**:3159-3176.
 16. Hohnjec N, Vieweg MF, Pühler A, Becker A, Küster H: **Overlaps in the transcriptional profiles of *Medicago truncatula* roots inoculated with two different *Glomus* fungi provide insights into the genetic program activated during arbuscular mycorrhiza.** *Plant Physiol* 2005, **137(4)**:1283-1301.
 17. Holmes P, Goffard N, Weiller GF, Rolfe BG, Imin N: **Transcriptional profiling of *Medicago truncatula* meristematic root cells.** *BMC Plant Biol* 2008, **8(1)**:21.
 18. Horak CE, Snyder M: **Global analysis of gene expression in yeast.** *Funct Integr Genomics* 2002, **2**:171-180.
 19. Czechowski T, Bari RP, Stitt M, Scheible WR, Udvardi MK: **Real-time RT-PCR profiling of over 1400 Arabidopsis transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific genes.** *Plant Journal* 2004, **38(2)**:366-379.
 20. Caldana C, Scheible WR, Mueller-Roeber B, Ruzicic S: **A quantitative RT-PCR platform for high-throughput expression profiling of 2500 rice transcription factors.** *Plant Methods* 2007, **3**:7.
 21. Scheible WR, Morcuende R, Czechowski T, Fritz C, Osuna D, Palacios-Rojas N, Schindelasch D, Thimm O, Udvardi MK, Stitt M: **Genome-wide reprogramming of primary and secondary metabolism, protein synthesis, cellular growth processes, and the regulatory infrastructure of Arabidopsis in response to nitrogen.** *Plant Physiology* 2004, **136(1)**:2483-2499.
 22. McGrath KC, Dombrecht B, Manners JM, Schenk PM, Edgar CI, Maclean DJ, Scheible WR, Udvardi MK, Kazan K: **Repressor- and activator-type ethylene response factors functioning in jasmonate signaling and disease resistance identified via a genome-wide screen of Arabidopsis transcription factor gene expression.** *Plant Physiol* 2005, **139(2)**:949-959.
 23. Morcuende R, Bari R, Gibon Y, Zheng W, Pant BD, Blasing O, Usadel B, Czechowski T, Udvardi MK, Stitt M, Scheible WR: **Genome-wide reprogramming of metabolism and regulatory networks of Arabidopsis in response to phosphorus.** *Plant Cell Environ* 2007, **30(1)**:85-112.
 24. Libault M, Wan J, Czechowski T, Udvardi M, Stacey G: **Identification of 118 Arabidopsis transcription factor and 30 ubiquitin-ligase genes responding to chitin, a plant-defense elicitor.** *Mol Plant Microbe Interact* 2007, **20(8)**:900-911.
 25. Guo A, He K, Liu D, Bai S, Gu X, Wei L, Luo J: **DATF: a database of Arabidopsis transcription factors.** *Bioinformatics* 2005, **21(10)**:2568-2569.
 26. Iida K, Seki M, Sakurai T, Satou M, Akiyama K, Toyoda T, Konagaya A, Shinozaki K: **RARTF: Database and tools for complete sets of Arabidopsis transcription factors.** *DNA Res* 2005, **12(4)**:247-256.
 27. **InterPro: Database of protein families, domains and sites** [<http://www.ebi.ac.uk/interpro/>]
 28. **WU-BLAST: Washington University BLAST Archives** [<http://blast.wustl.edu/>]
 29. **NCBI: The National Center for Biotechnology Information** [<http://www.ncbi.nlm.nih.gov/blast/>]
 30. **UniProt: The Universal Protein Resource** [<http://www.uniprot.org/>]
 31. Ramakers C, Ruijter JM, Deprez RH, Moorman AF: **Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data.** *Neurosci Lett* 2003, **339**:62-66.
 32. Pay A, Heberle-Bors E, Hirt H: **An alfalfa cDNA encodes a protein with homology to translationally controlled human tumor protein.** *Plant Molecular Biology* 1992, **19**:501-503.
 33. Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR: **Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis.** *Plant Physiology* 2005, **139**:5-17.
 34. **geNorm software** [<http://medgen.ugent.be/%7Ejvdesomp/genorm/>]
 35. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F: **Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes.** *Genome Biology* 2002, **3**:research0034.0031-research0034.0011.
 36. Chomczynski P, Mackey K: **Modification of the TRI reagent procedure for isolation of RNA from polysaccharide- and proteoglycan-rich sources.** *Biotechniques* 1995, **19(6)**:942-945.
 37. Rozen S, Skaletsky H: **Primer 3 on the WWW for general users and for biologist programmers.** In *Bioinformatics Methods and Protocols: Methods in Molecular Biology* Edited by: Krawetz S, Misener S. Totowa, NJ, Humana Press; 2000:365-386.
 38. Benedetto VA, Torres-Jerez I, Murray JD, Andriankaja A, Allen S, Kakar K, Wandrey M, Verdier J, Zuber H, Ott T, Moreau S, Niebel A, Frickey T, Weiller G, He J, Dai X, Zhao PX, Tang Y, Udvardi MK: **A gene expression atlas of the model legume *Medicago truncatula*.** *Plant Journal* 2008, (in press):.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

